

Gaucho® insecticidal seed treatment helps you optimise the productivity of your pasture and your profits and gives exceptional convenience and peace of mind. From the moment of sowing and throughout germination and early growth, Gaucho protects each and every seed and seedling from redlegged

earth mites and blue oat mites. Gaucho reduces
your need to spray, so that earthworms and
other beneficial species can flourish. As a
part of the Seedtech™ System by Bayer, Gaucho is

applied professionally and delivered on seed. Just order your Gaucho together with your pasture seed.







## An innovative seed treatment for pasture.

Gaucho® Flowable Seed Dressing offers farmers a new way to protect pasture seedlings against earth mite attack. The discovery of Gaucho (imidacloprid) by Bayer represented a true breakthrough in seed treatment technology. Gaucho is from the chloronicotinyl group of insecticides – an insecticide group with a unique mode of action.

Gaucho is highly systemic, making it an ideal seed treatment. It is rapidly absorbed by the seed and is translocated to the cotyledons and leaves. It has strong anti-feeding action against mites. The effect will last for up to 4 weeks and protects the pasture seedlings through the danger period of emergence and establishment.

Gaucho should be used as part of an integrated approach to mite control. The four main steps are:

- 1. A spring spray with Le-mat® 290 SL Insecticide to reduce adult mite numbers,
- 2. Use Gaucho as a seed treatment to repel earth mites from establishing plants,
- 3 A perimeter spray of Le-mat around the newly sown paddock to prevent earth mite invasions from neighbouring paddocks,
- 4. Monitor crops for re-invasion and if necessary apply Le-mat to reduce mite numbers.

Gaucho reduces the need for bare earth insecticides after planting. Gaucho has a low toxicity and is safe to beneficial species such as spiders, ants and ladybirds when used as directed. It is safe to soil microfauna and earthworms, and does not leach.




Figure 1. Plants/m<sup>2</sup> Gaucho vs untreated 4 weeks after sowing





