

# Grape Botrytis & Powdery Mildew positioning guide for non-export crops



## Seasonal Spray Timings

	E-L 1	E-L 3	E-L 9	E-L 12	E-L 20	E-L 25	E-L 29	E-L 35	E-L 37	E-L 43	E-L 47	
	Pruning	Woolly bud	2-3 leaves separated	5 leaves separated	Flowering 10% cap-fall	80% cap-fall	4 mm berries	Veraison	Pre-harvest	Post-harvest	Leaf fall	Dormancy
<b>Botrytis Domestic (non-export)</b>					Luna® Experience (to 4 mm berries (E-L 29))	Teldor® (to 21 days before harvest) Scala® (to 7 days before harvest)	Luna® Experience (to 4 mm berries (E-L 29)) Serenade® OPTI (No withholding period)	Group 9 + 12 (to 4 weeks before harvest) Teldor® (to 21 days before harvest) Scala® (to 7 days before harvest) Serenade® OPTI (No withholding period)		Luna® EXPERIENCE TEL DOR® SCALA® SERENADE® OPTI	GROUP 3 7 FUNGICIDE GROUP 17 FUNGICIDE GROUP 9 FUNGICIDE GROUP BM02 FUNGICIDE	
<b>Powdery Mildew Domestic (non-export)</b>	Grower Program		Prosper® (to 4 weeks before harvest)		Luna® Experience (to 4 mm berries (E-L 29))	Flint® (to 5 weeks before harvest)	Luna® Experience (to 4 mm berries (E-L 29))	Grower Program		Luna® EXPERIENCE PROSPER® FLINT®	GROUP 3 7 FUNGICIDE GROUP 5 FUNGICIDE GROUP 11 FUNGICIDE	

## Croplife Resistance Management Spray Guides

### Botrytis Resistance

Number of Botrytis sprays per season	Maximum recommended number of sprays which can contain Group:					
	2	7 (inc 7+3, 7+12)	9 (inc 9+12)	11 (inc 11+3)	12 (inc 7+12, 9+12)	17
1	1	1	1	1	1	1
2	1	1	1	1	2	1
3	1	1	1	1	2	1
4	2	2	2	2	2	2
5+	2	2	2	2	2	2

### Powdery Mildew Resistance

	Group						
	3	5	7	11	13	U6	50
Maximum number of consecutive sprays when applied as the only fungicide	2	2	1	1	2	2	2
Maximum number of consecutive sprays when applied as a tank-mix or co-formulant	2	2	2	2	2	2	2
Maximum number of sprays per season	3	3	3	2	3	2	4